



# Your fight against advanced colorectal cancer (MSI-H/dMMR) may start here

## Indication

OPDIVO® (nivolumab) is a prescription medicine that may be used in combination with YERVOY® (ipilimumab) to treat adults and children 12 years of age and older, with a type of colon or rectal cancer (colorectal cancer) that cannot be removed with surgery, or has spread, and is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR).

## How can OPDIVO + YERVOY help you fight advanced CRC (MSI-H/dMMR)?



- OPDIVO and YERVOY are immunotherapies, which work differently than chemotherapy
- Immunotherapies are treatments that work with the body's immune system to help fight disease

CRC=colorectal cancer; MSI-H=microsatellite instability-high; dMMR=mismatch repair deficient.

[Learn more](#) to see if OPDIVO + YERVOY immunotherapies could be right for you.

## Summary of Warnings and Precautions

OPDIVO® (nivolumab) and YERVOY® (ipilimumab) can cause problems that can sometimes become serious or life-threatening and can lead to death. Serious side effects may include lung problems; intestinal problems; liver problems; hormone gland problems; kidney problems; skin problems; eye problems; problems in other organs and tissues; and complications of stem cell transplant, including graft-versus-host disease (GVHD), that uses donor stem cells (allogeneic). Call or see your healthcare provider right away for any new or worsening signs or symptoms. OPDIVO is associated with severe infusion reactions.



Please see Important Safety Information about [OPDIVO and YERVOY](#), including serious side effects (page 7). For more information, please see [U.S. Full Prescribing Information](#) and [Medication Guide](#) for OPDIVO and [U.S. Full Prescribing Information](#) and [Medication Guide](#) for YERVOY, or talk to your healthcare team.



# Table of Contents

<a href="#">Introduction to OPDIVO + YERVOY.....</a>	<a href="#">1</a>
<a href="#">Could OPDIVO + YERVOY be right for you?.....</a>	<a href="#">3</a>
<a href="#">Clinical Trial Results: OPDIVO + YERVOY combination.....</a>	<a href="#">4</a>
<a href="#">Recognizing and managing side effects.....</a>	<a href="#">5</a>
<a href="#">Questions to ask your doctor.....</a>	<a href="#">6</a>
<a href="#">Important Safety Information about OPDIVO + YERVOY.....</a>	<a href="#">7-9</a>
<a href="#">Important Safety Information about OPDIVO QVANTIG.....</a>	<a href="#">9-12</a>
<a href="#">OPDIVO + YERVOY summary.....</a>	<a href="#">13</a>

## Could OPDIVO<sup>®</sup> + YERVOY<sup>®</sup> be effective against your cancer?

Knowing whether your colorectal cancer is **microsatellite instability-high (MSI-H)** or **mismatch repair deficient (dMMR)** can help you and your doctor choose the treatment that's right for you.

Ask your doctor if you have been tested for MSI-H or dMMR. In some cases, this testing is already done when you are first diagnosed, and your doctor can review your records to confirm your results.

### What are biomarkers?



Biomarkers help doctors select the appropriate treatment for you, so knowing your biomarker results can influence treatment decisions. Biomarkers are a protein, gene, or other substance in the body that reveals information about your tumor and how medicines may fight it.

Two biomarkers found in CRC that can inform the right treatment for you are MSI-H and dMMR.

### About the clinical trial

The clinical trial included 839 people with advanced CRC (MSI-H/dMMR) and compared OPDIVO + YERVOY to either a chemotherapy chosen by the investigators or OPDIVO alone.

In the one group, 354 people were given OPDIVO + YERVOY and 353 people were given OPDIVO alone.

In the other group, 202 people were given OPDIVO + YERVOY and 101 people were given investigator's choice of chemotherapy. Investigators in this trial were able to choose between giving modified FOLFOX-6 (leucovorin, fluorouracil, and oxaliplatin) with or without targeted therapy (bevacizumab or cetuximab) or FOLFIRI (irinotecan, leucovorin, and fluorouracil) with or without targeted therapy (bevacizumab or cetuximab).

### Summary of Warnings and Precautions

OPDIVO<sup>®</sup> (nivolumab) and YERVOY<sup>®</sup> (ipilimumab) can cause problems that can sometimes become serious or life-threatening and can lead to death. Serious side effects may include lung problems; intestinal problems; liver problems; hormone gland problems; kidney problems; skin problems; eye problems; problems in other organs and tissues; and complications of stem cell transplant, including graft-versus-host disease (GVHD), that uses donor stem cells (allogeneic). Call or see your healthcare provider right away for any new or worsening signs or symptoms. OPDIVO is associated with severe infusion reactions.

# OPDIVO + YERVOY is the only FDA-approved combination of 2 immunotherapies to treat advanced CRC (MSI-H/dMMR)

People given OPDIVO + YERVOY lived significantly longer without their cancer spreading, growing, or getting worse compared to OPDIVO alone at **~4 years**

## OPDIVO + YERVOY

### Not Reached

More than half the people went without their cancer spreading, growing, or getting worse

## OPDIVO Alone

### at 39 months

Half the people went without their cancer spreading, growing, or getting worse



**38% lower** risk of cancer spreading, growing, or getting worse compared to those given OPDIVO alone.

People given OPDIVO + YERVOY lived significantly longer without their cancer spreading, growing, or getting worse compared to investigator's choice chemotherapy at **~2.5 years**

## OPDIVO + YERVOY

### Not Reached

More than half the people went without their cancer spreading, growing, or getting worse

## Investigator's choice chemotherapy

### at 6 months

Half the people went without their cancer spreading, growing, or getting worse



**79% lower** risk of cancer spreading, growing, or getting worse compared to those given chemotherapy.

### Summary of Warnings and Precautions

OPDIVO® (nivolumab) and YERVOY® (ipilimumab) can cause problems that can sometimes become serious or life-threatening and can lead to death. Serious side effects may include lung problems; intestinal problems; liver problems; hormone gland problems; kidney problems; skin problems; eye problems; problems in other organs and tissues; and complications of stem cell transplant, including graft-versus-host disease (GVHD), that uses donor stem cells (allogeneic). Call or see your healthcare provider right away for any new or worsening signs or symptoms. OPDIVO is associated with severe infusion reactions.

## Recognizing and managing side effects

Although not every person may experience them, side effects are still possible with OPDIVO® + YERVOY®. Learn more about the most common side effects, potential side effects, and how to manage these side effects as well.

Please see important facts about potential side effects starting on page 6.

### Keep in mind:

- You may experience side effects during and after treatment
- Talk with your doctor if you notice anything unusual
- Informing your doctor about side effects can help you work together to manage your treatment plan



## [Learn more about potential side effects](#)

### Summary of Warnings and Precautions

OPDIVO® (nivolumab) and YERVOY® (ipilimumab) can cause problems that can sometimes become serious or life-threatening and can lead to death. Serious side effects may include lung problems; intestinal problems; liver problems; hormone gland problems; kidney problems; skin problems; eye problems; problems in other organs and tissues; and complications of stem cell transplant, including graft-versus-host disease (GVHD), that uses donor stem cells (allogeneic). Call or see your healthcare provider right away for any new or worsening signs or symptoms. OPDIVO is associated with severe infusion reactions.



# Questions to ask your doctor

Talking with your healthcare provider is a great place to start.

If you're considering OPDIVO + YERVOY as a treatment option for advanced CRC, these questions can help you start a conversation with your doctor at your next appointment.

Be sure to discuss your existing health problems, prescription medications, and non-prescription medicines, including vitamins and herbal supplements. Your doctor is always the best source of information when it comes to your health.

- What is immunotherapy?
- How is OPDIVO + YERVOY different from chemotherapy?
- How does OPDIVO + YERVOY work?
- What are the potential benefits of OPDIVO + YERVOY?
- How is OPDIVO + YERVOY given?
- Is an intravenous (IV) infusion my only option to receive treatment?
- How often will I need OPDIVO + YERVOY treatments, and how long will they take?
- What are the potential side effects that I may experience?
- Will this treatment affect my daily life in any way?
- How will I know if the treatment is working?
- What financial support is available?

Use this section to write down notes and questions you may have about OPDIVO + YERVOY.

---

---

---

---

---

---

---

---

---

---

## Summary of Warnings and Precautions

OPDIVO® (nivolumab) and YERVOY® (ipilimumab) can cause problems that can sometimes become serious or life-threatening and can lead to death. Serious side effects may include lung problems; intestinal problems; liver problems; hormone gland problems; kidney problems; skin problems; eye problems; problems in other organs and tissues; and complications of stem cell transplant, including graft-versus-host disease (GVHD), that uses donor stem cells (allogeneic). Call or see your healthcare provider right away for any new or worsening signs or symptoms. OPDIVO is associated with severe infusion reactions.

## Important Facts About OPDIVO® (nivolumab) + YERVOY® (ipilimumab)

This is a summary of important information that you need to know about OPDIVO + YERVOY. Your healthcare team can work with you to help answer any questions you may have about these medications. Keep this information in a safe place so you can refer to it before and during your treatment.

### Look out for the following icons as you read:



Talk to your healthcare team



Call a healthcare provider right away



Helpful information to remember

### What is OPDIVO + YERVOY?



OPDIVO is a prescription medicine used in combination with YERVOY (OPDIVO + YERVOY) to treat **colorectal cancer (a type of colon or rectal cancer)**. OPDIVO + YERVOY is for adults and children 12 years of age and older when:

- ✓ You have colorectal cancer that cannot be removed with surgery, or has spread to other parts of the body (metastatic), **AND**
- ✓ You have a tumor that is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR).

It is not known if OPDIVO is safe and effective in children younger than 12 years of age with MSI-H or dMMR metastatic colorectal cancer.

### What is the most important information I should know about OPDIVO + YERVOY?

OPDIVO and YERVOY are medicines that may treat certain cancers by working with your immune system. OPDIVO and YERVOY can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. Some of these problems may happen more often when OPDIVO is used in combination with another therapy.



**Get medical help immediately** if you develop any of these signs or symptoms or they get worse. It may keep these problems from becoming more serious. Your healthcare team will check you for these problems during treatment and may treat you with corticosteroid or hormone replacement medicines. If you have severe side effects, your healthcare team may also need to delay or completely stop your treatment.

### What are the serious side effects of OPDIVO + YERVOY?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening, and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

### Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

#### Lung problems – Things to look out for may include:

- cough
- shortness of breath
- chest pain

#### Intestinal problems – Things to look out for may include:

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdominal) pain or tenderness

#### Liver problems – Things to look out for may include:

- yellowing of your skin or the whites of your eyes
- pain on the right side of your stomach area (abdomen)
- bleeding or bruising more easily than normal
- severe nausea or vomiting
- dark urine (tea colored)

#### Hormone gland problems – Things to look out for may include:

- headaches that will not go away or unusual headaches
- weight gain or weight loss
- your voice gets deeper
- eye sensitivity to light
- feeling more hungry or thirsty than usual
- dizziness or fainting
- eye problems
- urinating more often than usual
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness
- rapid heartbeat
- hair loss
- feeling cold
- increased sweating
- constipation
- extreme tiredness

#### Kidney problems – Things to look out for may include:

- decrease in your amount of urine
- swelling of your ankles
- loss of appetite
- blood in your urine

#### Skin problems – Things to look out for may include:

- rash
- swollen lymph nodes
- fever or flu-like symptoms
- itching
- painful sore or ulcers in mouth or nose, throat, or genital area
- skin blistering or peeling

## Important Facts About OPDIVO + YERVOY

**Eye problems** – Things to look out for may include:

- blurry vision, double vision, or other vision problems
- eye pain or redness

**Problems can also happen in other organs and tissues. These are not all the signs and symptoms of immune system problems that can happen with OPDIVO and YERVOY. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:**

- Chest pain, irregular heartbeat, shortness of breath, swelling of ankles
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells, bruising

**Rejection of a transplanted organ or tissue.** Your healthcare provider should tell you what signs and symptoms you should report and monitor you depending on the type of organ transplant that you have had.

### What are the possible side effects of OPDIVO + YERVOY?

**OPDIVO + YERVOY can cause serious side effects, including:**

See the previous section, **“What is the most important information I should know about OPDIVO + YERVOY?”**

**Severe infusion reactions** – Things to look out for may include:

- chills or shaking
- shortness of breath or wheezing
- feel like passing out
- itching or rash
- fever
- flushing
- dizziness
- back or neck pain

**Tell your healthcare team** right away if you get these symptoms during an infusion of OPDIVO or YERVOY.

**Complications, including graft-versus-host disease (GVHD), of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).** These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with OPDIVO or YERVOY. Your healthcare provider will monitor you for these complications.

### What are the most common side effects?

The most common side effects of OPDIVO when used in combination with YERVOY include:

- feeling tired
- decreased appetite
- headache
- diarrhea
- vomiting
- low thyroid hormone levels (hypothyroidism)
- rash
- stomach-area (abdominal) pain
- constipation
- itching
- shortness of breath
- decreased weight
- nausea
- upper respiratory tract infection
- pain in muscles, bones, and joints
- fever
- cough

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

### What should I discuss with my healthcare team before receiving OPDIVO or YERVOY?

**Tell your healthcare provider about all your medical conditions, including if you:**

- have immune system problems such as Crohn’s disease, ulcerative colitis, or lupus
- have received an organ transplant, including corneal transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area in the past and have received other medicines that are like OPDIVO
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. OPDIVO and YERVOY can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if OPDIVO or YERVOY passes into your breast milk. Do not breastfeed during treatment and for 5 months after your last dose of OPDIVO or YERVOY.

## Important Facts About OPDIVO® (nivolumab) + YERVOY® (ipilimumab)



### Females who are able to become pregnant:

Your healthcare provider should do a pregnancy test before you start receiving OPDIVO or YERVOY.

- You should use an effective method of birth control during your treatment and for 5 months after the last dose of OPDIVO or YERVOY. Talk to your healthcare provider about birth control methods that you can use during this time.
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with OPDIVO or YERVOY. You or your healthcare provider should contact Bristol Myers Squibb at 1-844-593-7869 as soon as you become aware of a pregnancy.



### Tell your healthcare provider about all the medicines you take, including:

- prescription medicines
- over-the-counter medicines
- vitamins
- herbal supplements



For more information, please see accompanying U.S. Full Prescribing Information and Medication Guide for OPDIVO and for YERVOY, or talk to your healthcare team.

## Important Facts About OPDIVO QVANTIG™ (nivolumab + hyaluronidase-nvhy)

This is a summary of important information that you need to know about OPDIVO QVANTIG. Your healthcare team can work with you to help answer any questions you may have about this medication. Keep this document in a safe place, so you can refer to it before and during your treatment.

### Look out for the following icons as you read:



Talk to your healthcare team



Call a healthcare provider right away



Helpful information to remember

## What is OPDIVO QVANTIG?



**OPDIVO QVANTIG is a prescription medicine given as a subcutaneous injection (under the skin) used to treat adults with:**



A type of **colon or rectal cancer (colorectal cancer [CRC]):**

OPDIVO QVANTIG may be used alone, after completing combination treatment with intravenous nivolumab and ipilimumab, when colon or rectal cancer:

- ✓ Cannot be removed with surgery or has spread, **AND**
- ✓ Is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR)



It is not known if OPDIVO QVANTIG is safe and effective in children.



OPDIVO QVANTIG may not be used in combination with ipilimumab.

## Important Facts About OPDIVO QVANTIG

### What is the most important information I should know about OPDIVO QVANTIG?

OPDIVO QVANTIG is a medicine that may treat certain cancers by working with your immune system. OPDIVO QVANTIG can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. Some of these problems may happen more often when OPDIVO QVANTIG is used in combination with another therapy.

 **Get medical help immediately if you develop any of these signs or symptoms or they get worse.** It may keep these problems from becoming more serious. Your healthcare team will check you for these problems during treatment and may treat you with corticosteroid or hormone replacement medicines. If you have severe side effects, your healthcare team may also need to delay or completely stop your treatment.

### What are the serious side effects of OPDIVO QVANTIG?

A serious side effect is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

**Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms, including:**

**Lung problems** – Things to look out for may include:

- cough
- shortness of breath
- chest pain

**Intestinal problems** – Things to look out for may include:

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdominal) pain or tenderness

**Liver problems** – Things to look out for may include:

- yellowing of your skin or the whites of your eyes
- pain on the right side of your stomach area (abdomen)
- bleeding or bruising more easily than normal
- severe nausea or vomiting
- dark urine (tea colored)

**Hormone gland problems** – Things to look out for may include:

- headaches that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper
- dizziness or fainting
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

**Kidney problems** – Things to look out for may include:

- decrease in your amount of urine
- blood in your urine
- swelling of your ankles
- loss of appetite

**Skin problems** – Things to look out for may include:

- rash
- itching
- skin blistering or peeling
- swollen lymph nodes
- painful sores or ulcers in the mouth or nose, throat, or genital area
- fever or flu-like symptoms

**Problems can also happen in other organs and tissues.** These are not all of the signs and symptoms of immune system problems that can happen with OPDIVO QVANTIG.

 **Call or see your healthcare provider right away** for any new or worsening signs or symptoms, which may include:

- Chest pain, irregular heartbeat, shortness of breath, or swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells or bruising

**Rejection of a transplanted organ or tissue.** Your healthcare provider should tell you what signs and symptoms you should report and monitor you depending on the type of organ or tissue transplant that you have had.

## Important Facts About OPDIVO QVANTIG<sup>™</sup> (nivolumab + hyaluronidase-nvhy)

**Complications, including graft-versus-host disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).** These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with OPDIVO QVANTIG. Your healthcare provider will monitor you for signs of complications if you have an allogeneic stem cell transplant.

### What are the most common side effects of OPDIVO QVANTIG?

The most common side effects of OPDIVO QVANTIG when used alone in people with renal cell carcinoma include:

- pain in muscles, bones, and joints
- itchy skin
- diarrhea
- feeling tired
- rash
- cough
- low thyroid hormone levels
- stomach-area (abdominal) pain

The most common side effects observed with nivolumab given into the vein (intravenous nivolumab), which may be experienced with OPDIVO QVANTIG, are shown below.

The most common side effects of intravenous nivolumab when used alone include:

- feeling tired
- weakness
- upper respiratory tract infection
- rash
- cough
- fever
- pain in muscles, bones, and joints
- shortness of breath
- headache
- itching
- constipation
- stomach-area (abdominal) pain
- diarrhea
- decreased appetite
- vomiting
- nausea
- back pain
- urinary tract infection

The most common side effects of intravenous nivolumab when used in combination with cabozantinib as the first treatment for advanced RCC include:

- diarrhea
- rash, redness, pain, swelling, or blisters on the palms of your hands or soles of your feet
- pain in muscles, bones, and joints
- feeling tired
- liver problems. See “What is the most important information I should know about OPDIVO QVANTIG?”
- mouth sores
- decreased appetite
- nausea
- change in sense of taste
- high blood pressure
- rash
- stomach-area (abdominal) pain
- cough
- upper respiratory tract infection
- low thyroid hormone levels

The most common side effects of intravenous nivolumab when used in combination with platinum-containing chemotherapy and another chemotherapy medicine before having surgery for NSCLC include:

- nausea
- feeling tired
- rash
- constipation
- decreased appetite

The most common side effects of intravenous nivolumab when used in combination with cisplatin and gemcitabine to treat urothelial cancer include:

- nausea
- constipation
- vomiting
- feeling tired
- decreased appetite
- numbness, pain, tingling, or burning in your hands or feet
- pain in muscles, bones, and joints
- rash

The most common side effects of intravenous nivolumab when used in combination with fluoropyrimidine and platinum-containing chemotherapy to treat esophageal cancer and gastric cancer include:

- nausea
- decreased appetite
- diarrhea
- numbness, pain, tingling, or burning in your hands or feet
- feeling tired
- vomiting
- constipation
- stomach-area (abdominal) pain
- mouth sores
- pain in muscles, bones, and joints

These are not all the possible side effects of OPDIVO QVANTIG.

**Talk to your healthcare team** for more information about side effects. You are encouraged to report side effects of prescription drugs to the FDA by visiting [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or calling 1-800-FDA-1088.

### What should I discuss with my healthcare team before receiving OPDIVO QVANTIG?

**Talk to your healthcare team about all of your medical conditions,** including if you:

- have immune system problems such as Crohn’s disease, ulcerative colitis, or lupus
- have received an organ or tissue transplant, including corneal transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area in the past and have received other medicines that are like OPDIVO QVANTIG
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome



## Important Facts About OPDIVO QVANTIG

 **Talk to your healthcare team about all the medicines you are taking**, including:

- prescription medicines
- over-the-counter medicines
- vitamins
- herbal supplements

These are not all the topics you should discuss with your healthcare team.

Talk to your healthcare team about anything you might be unsure of before starting treatment.

### What should I discuss with my healthcare team about pregnancy, birth control, and breastfeeding?

 **Talk to your healthcare team** if:

 **You are pregnant or plan to become pregnant:** OPDIVO QVANTIG can harm your unborn baby.

If you are able to become pregnant, your healthcare provider should do a pregnancy test **before** you start receiving OPDIVO QVANTIG.

You should use an effective method of birth control **during your treatment and for 5 months after** your last dose of OPDIVO QVANTIG. Talk to your healthcare provider about birth control methods that you can use during this time.

 **Call your healthcare provider right away** if you become pregnant or think you may be pregnant during treatment with OPDIVO QVANTIG.

 **You are breastfeeding or plan to breastfeed:** It is not known if OPDIVO QVANTIG passes into your breast milk. Do not breastfeed **during treatment and for 5 months after** your last dose of OPDIVO QVANTIG.

### How will I receive OPDIVO QVANTIG?

 **OPDIVO QVANTIG is given as an injection under the skin** by your healthcare provider. An injection under the skin is sometimes called a **subcutaneous injection**. The injection is usually given in the stomach area (abdomen) or thigh, over about 3-5 minutes.

There are 3 parts to an **OPDIVO QVANTIG** appointment:

**Before** your injection:



Your healthcare provider will do blood tests to check you for side effects.

Your healthcare provider may delay or stop treatment with OPDIVO QVANTIG, if you have severe side effects.

**Receiving** your injection:



Injection time is **3-5 minutes**, though actual time in the clinic may vary.

**After** your injection:



Your healthcare provider may monitor you for side effects after your injection.

For a type of kidney cancer called advanced renal cell carcinoma, your healthcare provider may also prescribe you cabozantinib. Take cabozantinib exactly as your healthcare provider tells you.



OPDIVO QVANTIG is usually given every 2, 3, or 4 weeks, depending on the dose you are receiving.

Your healthcare provider will decide how many treatments you need.

### What should I do if I miss an appointment?



**Call your healthcare provider right away** to reschedule your appointment.



For more information, please see the U.S. Full Prescribing Information and Medication Guide for OPDIVO QVANTIG. Talk to your healthcare provider for more information about this medication.

**OPDIVO**  
(nivolumab)



**YERVOY**  
(ipilimumab)

Ask your doctor if  
**OPDIVO® + YERVOY®** is the  
right choice to treat your  
advanced CRC (MSI-H/dMMR)



**LEARN MORE**, to see if **OPDIVO + YERVOY**  
immunotherapies could be right for you

Actor portrayals.

OPDIVO + YERVOY is not approved for people younger than 12 years of age.  
OPDIVO Qvantig is not approved for people younger than 12 years of age.

## Discover **OPDIVO Qvantig™**, an immunotherapy treatment given by injection that's faster\*

**OPDIVO Qvantig™**  
nivolumab + hyaluronidase-nvhy  
SUBCUTANEOUS INJECTION | 120 mg + 2,000 units / mL

### Indication

OPDIVO QVANTIG™ (nivolumab and hyaluronidase-nvhy) is a prescription medicine used alone to treat adults with a type of colon or rectal cancer (colorectal cancer or CRC), after completing combination treatment with intravenous nivolumab and ipilimumab, when your cancer cannot be removed with surgery, or has spread, and is microsatellite instability-high (MSI-H) or mismatch repair deficient (dMMR).

OPDIVO QVANTIG may not be used in combination with ipilimumab.

It is not known if OPDIVO QVANTIG is safe and effective in children.

**You and your healthcare provider may have a choice between under-the-skin injections (subcutaneous injection) with **OPDIVO Qvantig** and intravenous (IV) infusions with **OPDIVO****

OPDIVO Qvantig may let you receive your immunotherapy treatment quickly—in as little as 3 minutes\*—instead of a 30-minute intravenous (IV) infusion of OPDIVO into a vein.

\*OPDIVO Qvantig injections are given by a healthcare provider in 3-5 minutes. This does not account for all aspects of treatment and does not include observation time. Actual clinic time may vary.

### Summary of Warnings and Precautions

OPDIVO QVANTIG™ (nivolumab + hyaluronidase-nvhy), OPDIVO® (nivolumab), and YERVOY® (ipilimumab) can cause problems that can sometimes become serious or life-threatening and can lead to death. Serious side effects may include lung problems; intestinal problems; liver problems; hormone gland problems; kidney problems; skin problems; eye problems; problems in other organs and tissues; and complications of stem cell transplant, including graft-versus-host disease (GVHD), that uses donor stem cells (allogeneic). Call or see your healthcare provider right away for any new or worsening signs or symptoms. OPDIVO is associated with severe infusion reactions.

**i** Please see Important Safety Information about [OPDIVO and YERVOY](#), including serious side effects. For more information, please see [U.S. Full Prescribing Information and Medication Guide](#) for OPDIVO and [U.S. Full Prescribing Information and Medication Guide](#) for YERVOY, or talk to your healthcare team.

 Bristol Myers Squibb®