



For Certain Previously Untreated Adults with Advanced Gastric,
Gastroesophageal Junction, and Esophageal Adenocarcinomas



A CHANCE TO LOOK FORWARD

OPDIVO will not work for everyone. Individual results may vary.

Please see information about the clinical trials on pages 3, 5, and 6. Please see serious side effects on the right panel.

What is OPDIVO?

OPDIVO is a prescription medicine used to treat people with cancer of the stomach (**gastric cancer**), cancer where the esophagus joins the stomach (**gastroesophageal junction cancer**), and in people with cancer of the tube that connects the throat to the stomach (**esophageal adenocarcinoma**). OPDIVO may be used in combination with chemotherapy that contains fluoropyrimidine and platinum in people whose cancer:

- ✓ Cannot be removed with surgery **OR**
- ✓ Has spread to other parts of the body.

It is not known if OPDIVO is safe and effective in children younger than 18 years of age.

Information provided in this brochure is not a substitute for talking with your healthcare professional.
Your healthcare professional is the best source of information about your disease.

OPDIVO
®
(nivolumab)

INJECTION FOR INTRAVENOUS USE 10 mg/mL




Please see Important Facts about OPDIVO, including serious side effects, on pages 9-10 and accompanying U.S. Full Prescribing Information and Medication Guide.



The **first and only** immunotherapy plus chemotherapy combination **FDA-approved for 3 types** of advanced gastroesophageal adenocarcinomas

OPDIVO is a prescription medicine used to treat people with cancer of the stomach (gastric), cancer where the esophagus joins the stomach (gastroesophageal junction), and in people with esophageal adenocarcinoma. OPDIVO may be used in combination with chemotherapy that contains fluoropyrimidine and platinum when your cancer cannot be removed with surgery **or** has spread to other parts of the body.



Select Important Safety Information

OPDIVO is a medicine that may treat certain cancers by working with your immune system. OPDIVO can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. Some of these problems may happen more often when OPDIVO is used in combination with another therapy.

What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- lung problems
- intestinal problems
- liver problems
- hormone gland problems
- kidney problems
- skin problems
- severe infusion reactions
- complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic)

What are the most common side effects?

The most common side effects of OPDIVO when used alone include:

- feeling tired
- rash
- pain in muscles, bones, and joints
- itchy skin
- diarrhea
- nausea
- weakness
- cough
- vomiting
- shortness of breath
- constipation
- decreased appetite
- back pain
- upper respiratory tract infection
- fever
- headache
- stomach-area (abdominal) pain
- urinary tract infection

The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:

- numbness, pain, tingling, or burning in your hands or feet
- nausea
- feeling tired
- diarrhea
- vomiting
- decreased appetite
- stomach-area (abdominal) pain
- constipation
- pain in muscles, bones, and joints

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

A treatment for people regardless of their PD-L1 biomarker status



WHAT'S A BIOMARKER?

A **PD-L1 biomarker** is a molecule found in the body in tissues, blood, or fluid. Biomarkers may show if there is a disease or condition present. To see if you have the PD-L1 biomarker, your doctor may test you and use a calculation called a **combined positive score (CPS)**.

*Programmed death-ligand 1.

OPDIVO + chemotherapy[†] may be right for you

While your doctor may test you for the PD-L1 biomarker to help you consider your treatment options,


OPDIVO + chemotherapy[†] is FDA-approved to treat advanced gastric, GEJ, and esophageal adenocarcinomas regardless of your test results.

ABOUT THE CLINICAL TRIAL

The study consisted of 1,581 patients, regardless of their PD-L1 biomarker status.

All had **previously untreated** advanced gastric, gastroesophageal junction, and esophageal adenocarcinomas.

Of these, 955 patients had a **PD-L1 CPS** of ≥ 5 .



[†]Modified FOLFOX-6 (leucovorin, fluorouracil, and oxaliplatin) or CapeOX (capecitabine and oxaliplatin).

Ask your doctor if OPDIVO + chemotherapy is right for you

What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- lung problems
- intestinal problems
- liver problems
- hormone gland problems
- kidney problems
- skin problems
- severe infusion reactions
- complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic)

What are the most common side effects?

The most common side effects of OPDIVO when used alone include:

- | | | |
|--------------------------------------|-----------------------|-------------------------------------|
| • feeling tired | • weakness | • upper respiratory tract infection |
| • rash | • cough | • fever |
| • pain in muscles, bones, and joints | • vomiting | • headache |
| • itchy skin | • shortness of breath | • stomach-area (abdominal) pain |
| • diarrhea | • constipation | • urinary tract infection |
| • nausea | • decreased appetite | |
| | • back pain | |

The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:

- | | | |
|--|---------------------------------|--------------------------------------|
| • numbness, pain, tingling, or burning in your hands or feet | • diarrhea | • constipation |
| • nausea | • vomiting | • pain in muscles, bones, and joints |
| • feeling tired | • decreased appetite | |
| | • stomach-area (abdominal) pain | |

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

More people were alive on OPDIVO + chemotherapy* compared to chemotherapy* alone



In a study of 1,581 patients with previously untreated advanced gastric, gastroesophageal junction, and esophageal adenocarcinomas, 789 received OPDIVO in combination with chemotherapy* and 792 received chemotherapy* alone. Of these, 955 patients had a PD-L1 CPS of ≥ 5 . In this group, 473 patients received OPDIVO in combination with chemotherapy* while 482 patients received chemotherapy* alone.

Patients with PD-L1 CPS ≥ 5 (n=955)

PEOPLE ALIVE AT 1 YEAR

57%

compared to

46%

on chemotherapy*
alone

OPDIVO + chemotherapy*

proven to  lower the risk of dying by **29%**
compared to **chemotherapy* alone**

Half of the patients who were on **OPDIVO + chemotherapy*** were alive at 14.4 months compared to half being alive on **chemotherapy* alone** at 11.1 months.

All patients regardless of PD-L1 CPS status (n=1,581)

PEOPLE ALIVE AT 1 YEAR

55%

compared to

48%

on **OPDIVO + chemotherapy***

OPDIVO + chemotherapy*
proven to  lower the risk of dying by **20%**
compared to **chemotherapy* alone**

Half of the patients who were on **OPDIVO + chemotherapy*** were alive at 13.8 months compared to half being alive on **chemotherapy* alone** at 11.6 months.

For patients with a PD-L1 biomarker CPS of ≥ 5 , half of the patients on **OPDIVO + chemotherapy*** went 7.7 months without their cancer spreading, growing, or getting worse compared to half who went without their cancer spreading, growing, or getting worse on **chemotherapy* alone** at 6.0 months.

OPDIVO + chemotherapy* reduced the risk of cancer spreading, growing, or getting worse by 32% compared to **chemotherapy* alone**.

*Modified FOLFOX-6 (leucovorin, fluorouracil, and oxaliplatin) or CapeOX (capecitabine and oxaliplatin).

OPDIVO will not work for everyone. Individual results may vary.

What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- lung problems
- intestinal problems
- liver problems
- hormone gland problems
- kidney problems
- skin problems
- severe infusion reactions
- complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic)

What are the most common side effects?

The most common side effects of OPDIVO when used alone include:

- feeling tired
- rash
- pain in muscles, bones, and joints
- itchy skin
- diarrhea
- nausea
- weakness
- cough
- vomiting
- shortness of breath
- constipation
- decreased appetite
- back pain
- upper respiratory tract infection
- fever
- headache
- stomach-area (abdominal) pain
- urinary tract infection

The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:

- numbness, pain, tingling, or burning in your hands or feet
- nausea
- feeling tired
- diarrhea
- vomiting
- decreased appetite
- stomach-area (abdominal) pain
- constipation
- pain in muscles, bones, and joints

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

i Please see additional Important Facts about OPDIVO, including serious side effects, on pages 9-10 and accompanying U.S. Full Prescribing Information and Medication Guide.

More people had their tumors shrink or disappear on OPDIVO + chemotherapy* compared to chemotherapy* alone

In a study of 1,581 patients with previously untreated advanced gastric, gastroesophageal junction, and esophageal adenocarcinomas, 789 received **OPDIVO in combination with chemotherapy*** and 792 received **chemotherapy* alone**. Of these, 955 patients had a PD-L1 CPS of ≥ 5 . In this group, 473 patients received OPDIVO in combination with chemotherapy* while 482 patients received chemotherapy* alone.

Patients with PD-L1 CPS ≥ 5 (n=955)

	OPDIVO + chemotherapy* (n=473)	chemotherapy* alone (n=482)
Tumors disappeared completely (complete response) [†]	12% (n=55)	7% (n=34)
Tumors shrank (partial response)	38% (n=182)	31% (n=150)
Tumors shrank or disappeared completely (overall response)	50% (n=237)	38% (n=184)

All patients regardless of PD-L1 CPS status (n=1,581)

	OPDIVO + chemotherapy* (n=789)	chemotherapy* alone (n=792)
Tumors disappeared completely (complete response) [†]	10% (n=78)	7% (n=52)
Tumors shrank (partial response)	37% (n=292)	30% (n=241)
Tumors shrank or disappeared completely (overall response)	47% (n=370)	37% (n=293)

[†]The disappearance of any measurable tumors in response to treatment.
This does not necessarily mean the cancer has been cured.

[†]The disappearance of any measurable tumors in response to treatment.
This does not necessarily mean the cancer has been cured.

*Modified FOLFOX-6 (leucovorin, fluorouracil, and oxaliplatin) or CapeOX (capecitabine and oxaliplatin).

OPDIVO will not work for everyone. Individual results may vary.

What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- lung problems
- intestinal problems
- liver problems
- hormone gland problems
- kidney problems
- skin problems
- severe infusion reactions
- complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic)

What are the most common side effects?

The most common side effects of OPDIVO when used alone include:


- | | | |
|--------------------------------------|-----------------------|-------------------------------------|
| • feeling tired | • weakness | • upper respiratory tract infection |
| • rash | • cough | • fever |
| • pain in muscles, bones, and joints | • vomiting | • headache |
| • itchy skin | • shortness of breath | • stomach-area (abdominal) pain |
| • diarrhea | • constipation | • urinary tract infection |
| • nausea | • decreased appetite | |
| | • back pain | |

The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:

- | | | |
|--|---------------------------------|--------------------------------------|
| • numbness, pain, tingling, or burning in your hands or feet | • diarrhea | • constipation |
| • nausea | • vomiting | • pain in muscles, bones, and joints |
| • feeling tired | • decreased appetite | |
| | • stomach-area (abdominal) pain | |

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

OPDIVO Helps Seek and Attack: How it works with your immune system



OPDIVO can cause your T cells to attack healthy cells contained within organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become serious or life-threatening and can lead to death, and may happen anytime during treatment or even after treatment has ended. You may have more than one of these problems at the same time. Your doctor will check you for these problems during treatment. Your doctor may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may also need to delay or completely stop treatment if you have severe side effects. Please see pages 9-10 for a list of possible side effects and what to look out for.

What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- lung problems
- intestinal problems
- liver problems
- hormone gland problems
- kidney problems
- skin problems
- severe infusion reactions
- complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic)

What are the most common side effects?

The most common side effects of OPDIVO when used alone include:

- | | | |
|--------------------------------------|-----------------------|-------------------------------------|
| • feeling tired | • weakness | • upper respiratory tract infection |
| • rash | • cough | • fever |
| • pain in muscles, bones, and joints | • vomiting | • headache |
| • itchy skin | • shortness of breath | • stomach-area (abdominal) pain |
| • diarrhea | • constipation | • urinary tract infection |
| • nausea | • decreased appetite | |
| | • back pain | |

The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:

- | | | |
|--|---------------------------------|--------------------------------------|
| • numbness, pain, tingling, or burning in your hands or feet | • diarrhea | • constipation |
| • nausea | • vomiting | • pain in muscles, bones, and joints |
| • feeling tired | • decreased appetite | |
| | • stomach-area (abdominal) pain | |

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

OPDIVO + chemotherapy* is given every 2 or 3 weeks




OPDIVO + CHEMOTHERAPY* every 2 weeks

OPDIVO 240 mg + chemo*

OPDIVO is delivered into your vein through an intravenous (IV) line. You'll be given OPDIVO + chemotherapy* on the same day every 2 weeks.

Infusion time 30 minutes OPDIVO



CONTINUE TREATMENT


Treatment continues as your doctor monitors your condition to ensure treatment is working and side effects are manageable, for up to 2 years

OPDIVO + CHEMOTHERAPY* every 3 weeks

OPDIVO 360 mg + chemo*

OPDIVO is delivered into your vein through an intravenous (IV) line. You'll be given OPDIVO + chemotherapy* on the same day every 3 weeks.

Infusion time 30 minutes OPDIVO



CONTINUE TREATMENT

Treatment continues as your doctor monitors your condition to ensure treatment is working and side effects are manageable, for up to 2 years

*Your doctor will prescribe you a type of chemotherapy that contains fluoropyrimidine and platinum.

What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- lung problems
- intestinal problems
- liver problems
- hormone gland problems
- kidney problems
- skin problems
- severe infusion reactions
- complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic)

What are the most common side effects?

The most common side effects of OPDIVO when used alone include:

- feeling tired
- rash
- pain in muscles, bones, and joints
- itchy skin
- diarrhea
- nausea
- weakness
- cough
- vomiting
- shortness of breath
- constipation
- decreased appetite
- back pain
- upper respiratory tract infection
- fever
- headache
- stomach-area (abdominal) pain
- urinary tract infection

The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:

- numbness, pain, tingling, or burning in your hands or feet
- nausea
- feeling tired
- diarrhea
- vomiting
- decreased appetite
- stomach-area (abdominal) pain
- constipation
- pain in muscles, bones, and joints

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.



Your doctor will choose the chemotherapy option that's right for you



OPDIVO is given through intravenous (IV) infusion. The medicine is given directly into the bloodstream through a needle placed in a vein by a healthcare professional—usually in the arm or hand. Chemotherapy will be given on the same day.



Talk to your doctor to learn how many treatments may be right for you. Your doctor will decide how many treatments you need. Your healthcare provider will do blood tests to check you for side effects, and may treat you with corticosteroid or hormone replacement medicines, and delay or completely stop treatment if you have side effects that are severe. Be sure to talk to your healthcare team about side effects.

Frequent communication with your healthcare team is an important part of treatment. Use your appointments to let them know how you're feeling and ask any questions you may have. You may also be able to talk with others going through a similar situation.

See page 10 for information about pregnancy, nursing, health problems, and other medicines that you should discuss with your healthcare team before starting treatment.



Visit www.OPDIVO.com for more information.



If you miss an appointment, call a healthcare provider right away. Try to schedule your appointments for the same day of the week and the same time of day to make them easier to remember. It is important for you to keep all appointments with your healthcare provider.

What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- lung problems
- intestinal problems
- liver problems
- hormone gland problems
- kidney problems
- skin problems
- severe infusion reactions
- complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic)

What are the most common side effects?

The most common side effects of OPDIVO when used alone include:

- feeling tired
- rash
- pain in muscles, bones, and joints
- itchy skin
- diarrhea
- nausea
- weakness
- cough
- vomiting
- shortness of breath
- constipation
- decreased appetite
- back pain
- upper respiratory tract infection
- fever
- headache
- stomach-area (abdominal) pain
- urinary tract infection

The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:


- numbness, pain, tingling, or burning in your hands or feet
- nausea
- feeling tired
- diarrhea
- vomiting
- decreased appetite
- stomach-area (abdominal) pain
- constipation
- pain in muscles, bones, and joints

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.


Important Facts About OPDIVO® (nivolumab)

This is a summary of important information that you need to know about OPDIVO. Your healthcare team can work with you to help answer any questions you may have about OPDIVO. Keep this information in a safe place so you can refer to it before and during your treatment.


Look out for the following icons as you read:



Talk to your healthcare team



Call a healthcare provider right away



Helpful information to remember

What is OPDIVO?

OPDIVO is a prescription medicine used to treat people with cancer of the stomach (**gastric cancer**), cancer where the esophagus joins the stomach (**gastroesophageal junction cancer**), and in people with cancer of the tube that connects the throat to the stomach (**esophageal adenocarcinoma**). OPDIVO may be used in combination with chemotherapy that contains fluoropyrimidine and platinum in people whose cancer:

- ✓ Cannot be removed with surgery **OR**
- ✓ Has spread to other parts of the body.

It is not known if OPDIVO is safe and effective in children younger than 18 years of age.

What is the most important information I should know about OPDIVO?

OPDIVO is a medicine that may treat certain cancers by working with your immune system. OPDIVO can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. Some of these problems may happen more often when OPDIVO is used in combination with another therapy.



Get medical help immediately if you develop any of these signs or symptoms or they get worse. It may keep these problems from becoming more serious. Your healthcare team will check you for these problems during treatment and may treat you with corticosteroid or hormone replacement medicines. If you have severe side effects, your healthcare team may also need to delay or completely stop your treatment.

What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

Lung problems – Things to look out for may include:

- new or worsening cough
- shortness of breath
- chest pain

Intestinal problems – Things to look out for may include:

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdominal) pain or tenderness

Liver problems – Things to look out for may include:

- yellowing of your skin or the whites of your eyes
- severe nausea or vomiting
- pain on the right side of your stomach area (abdomen)
- dark urine (tea colored)
- bleeding or bruising more easily than normal

Hormone gland problems – Things to look out for may include:

- headaches that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper
- dizziness or fainting
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

Kidney problems – Things to look out for may include:

- decrease in your amount of urine
- blood in your urine
- swelling of your ankles
- loss of appetite

Skin problems – Things to look out for may include:

- rash
- itching
- skin blistering or peeling
- painful sores or ulcers in the mouth or nose, throat, or genital area

Problems can also happen in other organs and tissues. These are not all the signs and symptoms of immune system problems that can happen with OPDIVO. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:

- Chest pain, irregular heartbeat, shortness of breath, swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells, bruising



For more information, please see U.S. Full Prescribing Information and Medication Guide for OPDIVO, or talk to your healthcare team.

What are the possible side effects of OPDIVO?

OPDIVO can cause serious side effects, including:

See the previous section, “**What is the most important information I should know about OPDIVO?**”

Severe infusion reactions – Things to look out for may include:

- chills or shaking
- shortness of breath or wheezing
- feel like passing out
- itching or rash
- fever
- flushing
- dizziness
- back or neck pain

 **Tell your healthcare team** right away if you get these symptoms during an infusion of OPDIVO.

Complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic).

These complications can be severe and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with OPDIVO. Your healthcare provider will monitor you for these complications.

What are the most common side effects of OPDIVO?

The most common side effects of OPDIVO when used alone include:

- | | | |
|---|------------------------------|--|
| • feeling tired | • cough | • upper respiratory tract infection |
| • rash | • vomiting | • headache |
| • pain in muscles, bones, and joints | • shortness of breath | • stomach-area (abdominal) pain |
| • itchy skin | • constipation | • urinary tract infection |
| • diarrhea | • decreased appetite | |
| • nausea | • back pain | |
| • weakness | • fever | |

The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:

- | | | |
|---|--|---|
| • numbness, pain, tingling, or burning in your hands or feet | • diarrhea | • constipation |
| • nausea | • vomiting | • pain in muscles, bones, and joints |
| • feeling tired | • decreased appetite | |
| | • stomach-area (abdominal) pain | |

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

What should I discuss with my healthcare team before receiving OPDIVO?

 **Tell your healthcare provider about all your medical conditions, including if you:**

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area in the past and have received other medicines that are like OPDIVO
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. OPDIVO can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if OPDIVO passes into your breast milk. Do not breastfeed during treatment with OPDIVO and for 5 months after the last dose of OPDIVO.



Females who are able to become pregnant:

Your healthcare provider should do a pregnancy test before you start receiving OPDIVO.

- You should use an effective method of birth control during your treatment and for at least 5 months after the last dose of OPDIVO. Talk to your healthcare provider about birth control methods that you can use during this time.
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with OPDIVO.



Tell your healthcare provider about all the medicines you take, including:

- | | |
|------------------------------|----------------------|
| • prescription medicines | • vitamins |
| • over-the-counter medicines | • herbal supplements |



For more information, please see [U.S. Full Prescribing Information and Medication Guide for OPDIVO](#), or talk to your healthcare team.

Financial Resources

Patient Access, Reimbursement, and Co-Pay Support

Available through
Bristol Myers Squibb Access Support®

Bristol Myers Squibb is committed to helping patients gain access to their prescribed BMS medications. That's why we offer the BMS Access Support program, which provides resources to help patients understand their insurance coverage and find information on sources of financial support, including co-pay assistance for eligible commercially insured patients.

For more information, ask your doctor, visit **BMS Access Support**, or call BMS Access Support at **1-800-861-0048**, 8 AM to 8 PM ET, Monday–Friday.

Other Patient Resources*



CancerCare
cancercare.org

The mission of CancerCare is to provide free, professional support services to anyone affected by cancer.



Cancer Support Community
cancersupportcommunity.org

The Cancer Support Community is an international nonprofit dedicated to providing support, education, and hope to people affected by cancer and their loved ones.



Cancer Research Institute
cancerresearch.org

Cancer Research Institute (CRI) is dedicated to pursuing research in and advancing awareness of immunotherapy for cancer.



Debbie's Dream Foundation: Curing Stomach Cancer
debbiesdream.org

Debbie's Dream Foundation: Curing Stomach Cancer is a 501(c)(3) non-profit organization dedicated to raising awareness about stomach cancer, advancing funding for research, and providing education and support internationally to patients, families, and caregivers.



Esophageal Cancer Action Network (ECAN)
ECAN.org

ECAN is dedicated to saving lives by sounding the alarm about the link between reflux disease and cancer, promoting early detection, advocating for funding and progress in medical research, and educating patients and caregivers within a supportive, compassionate community.



EC Aware
ec-aware.org

The mission of EC Aware (formerly ECAA) is to promote awareness of the disease for prevention and to foster community that together can provide support and information to those in need.

*Bristol Myers Squibb does not endorse these organizations. The information and links provided by Bristol Myers Squibb are meant for informational purposes only and are not meant to replace a physician's medical advice.



For more information, please see [U.S. Full Prescribing Information and Medication Guide for OPDIVO](#), or talk to your healthcare team.



SIGN UP for OPDIVO with You.

OPDIVO with You means just that: we're here to help you better understand OPDIVO, whether you or a loved one are just considering OPDIVO or already have an OPDIVO prescription. This complimentary program includes the following resources. Sign up and you'll receive:



Helpful Information

Get useful tips, tools, and educational materials that can help you learn more about OPDIVO and may help support your care.



Support and Advocacy Groups

We can direct you to organizations that may be helpful for the challenges you may face.



A Care Counselor as Close as Your Phone

If you are prescribed OPDIVO and sign up for this additional resource, you'll be able to speak with a registered nurse. This personal Care Counselor can answer questions about treatment and help you to be a more informed partner with your doctor. Care Counselors cannot provide medical advice. Your healthcare professional is the best source of information about your health.

If you'd like to sign up for the **OPDIVO with You** program, or if you have general questions about OPDIVO, call **1-855-OPDIVO-1** (1-855-673-4861)

8 AM to 8 PM ET, Monday to Friday, or visit www.OPDIVOwithYou.com



For more information, call **1-855-OPDIVO-1** or visit www.OPDIVO.com

Bristol Myers Squibb is committed to helping patients throughout their treatment.



What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- | | | |
|-----------------------|-----------------------------|---|
| • lung problems | • hormone gland problems | • complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic) |
| • intestinal problems | • kidney problems | |
| • liver problems | • skin problems | |
| | • severe infusion reactions | |

What are the most common side effects?

The most common side effects of OPDIVO when used alone include:

- | | | |
|--------------------------------------|-----------------------|-------------------------------------|
| • feeling tired | • weakness | • upper respiratory tract infection |
| • rash | • cough | • fever |
| • pain in muscles, bones, and joints | • vomiting | • headache |
| • itchy skin | • shortness of breath | • stomach-area (abdominal) pain |
| • diarrhea | • constipation | • urinary tract infection |
| • nausea | • decreased appetite | |
| | • back pain | |

The most common side effects of OPDIVO when used in combination with fluoropyrimidine and platinum-containing chemotherapy include:

- | | | |
|--|---------------------------------|--------------------------------------|
| • numbness, pain, tingling, or burning in your hands or feet | • diarrhea | • constipation |
| • nausea | • vomiting | • pain in muscles, bones, and joints |
| • feeling tired | • decreased appetite | |
| | • stomach-area (abdominal) pain | |

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

i Please see additional Important Facts about OPDIVO, including serious side effects, on pages 9–10 and accompanying U.S. Full Prescribing Information and Medication Guide.



© 2021 Bristol-Myers Squibb Company.

OPDIVO®, Access Support®, A Chance to Live Longer™, and the related logos are trademarks of Bristol-Myers Squibb Company. All other trademarks listed are the property of their respective owners.

1506-US-2101684 10/21



INJECTION FOR INTRAVENOUS USE 10 mg/mL